


WYANDOT COUNTY 2016 COMMUNICABLE DISEASE REPORT

February 2017

Wyandot County saw a 0.87% increase in communicable disease cases from 2015 to 2016 (115 cases and 116 cases respectively).

Numerous infectious diseases were reported during 2016; however, the most frequently reported illnesses were chlamydia (43 cases), Hepatitis



C (19 cases), campylobacteriosis (18 cases), cryptosporidiosis (5 cases), and influenza-associated hospitalizations (5 cases). Chlamydia, Hepatitis C, campylobacteriosis, and influenza-associated hospitalizations were also in the top five diseases reported during 2014 and 2015. **Table 1.** on Page 2 illustrates all of the diseases reported in the community and the number of cases for each of these illnesses. The remainder of this document provides epidemiological data on each of these illnesses as well as brief demographic information on the cases and disease trends over the past five years.


Inside this issue:

Communicable Disease Table	2
Communicable Disease Graph	3
Chlamydia	4
Hepatitis C	5
Campylobacteriosis	6
Cryptosporidiosis	7
Influenza-Associated Hospitalizations	8
Timeliness of Reporting	9

Table 1. Communicable Diseases Reported in Wyandot County, 2016

Class B	
Campylobacteriosis	18
Chlamydia	43
Cryptosporidiosis	5
E. coli	1
Giardia	1
Gonorrhea	3
<i>Haemophilus influenzae</i>	1
Hepatitis B	4
Hepatitis C	19
Influenza-Associated Hospitalization	5
Legionnaires' Disease	1
Lyme Disease	2
Malaria	1
Meningitis - aseptic	1
Mycobacterial Disease	3
Salmonella	4
Shigella	1
<i>Streptococcal pneumoniae</i>	2
Varicella (Chicken Pox)	1
Total:	116

Types of Communicable Diseases Reported in Wyandot County, 2016



Notes:

Case counts include confirmed, probable, and suspect disease classifications

Sexually transmitted infections include chlamydia and gonorrhea

Enteric illnesses include campylobacteriosis, cryptosporidiosis, *E. coli*, giardia, salmonella, and shigella

Vaccine preventable illnesses include *Haemophilus influenzae*, Hepatitis B, influenza-associated hospitalizations, *Streptococcus pneumoniae*, and varicella

Bloodborne pathogens include Hepatitis C

Vectorborne illnesses include Lyme disease and Malaria

Other illnesses include Legionnaires' disease, aseptic meningitis, and mycobacterial disease

Chlamydia

Number of Cases: 43

Average Age: 23.8 years

Median Age: 21 years

Age Range: 13-43 years

Female: 76.7%

Male: 23.3%

Percent Change from 2015: 0.0%

EPIDEMIOLOGY

Infectious Agent: *Chlamydia trachomatis* bacteria

Case Definition: Isolation of *Chlamydia trachomatis* from a clinical specimen


Symptoms: Men may suffer from painful urination, urinary frequency, and penile discharge; while women may experience vaginal discharge

Source: Humans

Mode of Transmission: Sexually transmitted

Incubation Period: 7-21 days

Prevention: Abstinence, condom use, and identification and treatment of sexual contacts of those proven to be or suspected of being infected with *Chlamydia trachomatis*



Hepatitis C

Number of Cases: 19

Average Age: 33.6 years

Median Age: 29 years

Age Range: 18-61 years

Female: 63.2%

Male: 36.8%

Decrease from 2015: 9.5%

EPIDEMIOLOGY

Infectious Agent: Hepatitis C virus

Case Definition: A positive test for Hepatitis C virus antibodies or detection of the Hepatitis C virus


Symptoms: Those infected may be asymptomatic; however, some may experience nausea, vomiting, abdominal pain, loss of appetite, dark urine, and/or jaundice

Source: Human blood

Mode of Transmission: Injection drug use through the sharing of needles and other drug paraphernalia contaminated with infected blood; non-professional tattooing or in-home tattooing through shared needles or contaminated equipment; sexual transmission inefficiently spreads the virus (rare)

Incubation Period: 2 weeks – 6 months

Prevention: No vaccine is available



Campylobacteriosis

Number of Cases: 18

Average Age: 46.3 years

Median Age: 45 years

Age Range: 3-93 years

Female: 72.2%

Male: 27.8%

Increase from 2015: 12.5%

EPIDEMIOLOGY

Infectious Agent: *Campylobacter* organisms, most commonly *Campylobacter jejuni*

Case Definition: Isolation or detection of *Campylobacter* species with or without diarrhea


Symptoms: Diarrhea (frequently bloody), abdominal cramps, malaise, fever, headache, nausea, and vomiting

Source: Poultry, cattle, puppies, kittens, swine, sheep, rodents, and birds

Mode of Transmission: Fecal-orally through undercooked meat, contaminated food or raw milk, or direct contact with infected pets, livestock, or infants

Incubation Period: 1-10 days

Prevention: Cook meats thoroughly, avoid cross-contamination of foods with raw meat juices, avoid unpasteurized milk and untreated water, and washing hands after contact with animals, bowel movements, changing diapers, and before eating or preparing food



Cryptosporidiosis

Number of Cases: 5

Average Age: 15.6 years

Median Age: 7 years

Age Range: 13-65 years

Female: 60.0%

Male: 40.0%

Increase from 2015: 150.0%

EPIDEMIOLOGY

Infectious Agent: *Cryptosporidium* species

Case Definition: Evidence of *Cryptosporidium* organisms or DNA


Symptoms: Diarrhea, nausea, vomiting, abdominal cramping, fever, weight loss, and loss of appetite

Source: Humans, cattle (especially calves), and other domestic animals

Mode of Transmission: Occurs through the fecal-oral route, including person-to person, animal-to-person, waterborne, and foodborne (including raw milk and unpasteurized cider).

Incubation Period: 1-12 days

Prevention: Cook meats thoroughly; avoid unpasteurized milk, dairy products, juice and cider; wash hands thoroughly after handling household pets, farm animals or after working in soil; do not drink or swallow water from rivers, lakes, streams, the ocean, swimming pools, hot tubs or Jacuzzis; do not swim for two weeks after diarrhea ceases



Influenza -Associated Hospitalizations

Number of Cases: 5

Average Age: 42.0 years

Median Age: 55 years

Age Range: 13-65 years

Female: 80.0%

Male: 20.0%

Decrease from 2015: 16.7%

EPIDEMIOLOGY

Infectious Agent: Two main types of Influenza virus: Influenza A and Influenza B; both types include different strains that tend to change from year to year

Case Definition: An illness compatible with influenza virus infection that results in hospitalization


Symptoms: Fever, body aches, headache, malaise, nonproductive cough, sore throat, and runny nose

Source: Humans

Mode of Transmission: Direct person-to-person contact through droplet spread or via articles recently contaminated with nasopharyngeal secretions

Incubation Period: 1- 4 days

Prevention: The best prevention is annual vaccination; washing hands after sneezing, coughing, or using a tissue; cough into sleeve and not into hands



Timeliness of Disease Reporting

Timely reporting of infectious diseases is important in identifying potential outbreaks and in reducing disease burden. Public health relies on health care providers and laboratories for identification and prompt reporting of these infectious diseases. Timeliness requirements for each reportable disease is dependent on the infectious nature and severity of the disease. Reporting lag is defined as the difference between the date the case was reported to the local health department and the date of diagnosis. For Class A diseases, median and mean lag time values should be less than 1 since these illnesses are required to be reported to the health department immediately, and for Class B and C diseases, mean and median lag time values should be less than 2 since these illnesses should be reported to the health department by the end of the next business day. Table 2. illustrates the lag time for select Class B reportable diseases reported in Wyandot County during 2016.

Table 2. Reporting Lag Time for Select Reportable Diseases in Wyandot County, 2016

Reportable Disease	Reporting Requirement	Cases (N)	Median (Days)	Mean (Days)
Campylobacteriosis	By end of next business day	18	1	1.8
Cryptosporidiosis	By end of next business day	5	1	3
E. coli O157:H7	By end of next business day	1	21	21
Giardia	By end of next business day	1	3	3
Influenza-Associated Hospitalization	By end of next business day	5	2	3
Legionnaires' Disease	By end of next business day	1	5	5
Pertussis	By end of next business day	0	N/A	N/A
Salmonella	By end of next business day	4	1	1.3
Shigella	By end of next business day	1	0	0

Note: Reporting lag time is the difference between the date the case was reported to the local health department and the case's date of diagnosis
Date of diagnosis defaulted to lab specimen collection date or illness onset date if blank

Wyandot County



Public Health

Prevent. Promote. Protect.

**127-A S. Sandusky Street
Upper Sandusky, Ohio 43351
(419) 294-3852
<http://www.wyandotohealth.com>**

Compiled and prepared by:

Mary E. Salimbene Merriman,
MPH-Epidemiologist Zachary J.
Colles, MPH-Epidemiologist



**Division of Health Promotion and
Planning
940 London Avenue
Suite 1100
Marysville, Ohio 43040
(937) 642-2053**