



WYANDOT COUNTY 2017

COMMUNICABLE DISEASE REPORT

The communicable disease summary of reportable infectious disease for January 2017—December 2017.


TABLE OF CONTENTS

Annual Communicable Diseases	3
Communicable Disease Summary	3
Communicable Disease Table.....	4
Communicable Disease Graph.....	5
Chlamydia	6
Campylobacteriosis	7
Influenza-Associated Hospitalizations.....	8
Hepatitis C	9
Gonorrhea	10
Timeliness of Reporting.....	11

ANNUAL COMMUNICABLE DISEASES

COMMUNICABLE DISEASE SUMMARY

Wyandot County saw a 35.34% increase in communicable disease cases from 2016 to 2017 (116 cases and 157 cases, respectively).



Numerous infectious diseases were reported during 2017; however, the most frequently reported illnesses were chlamydia (57 cases), campylobacteriosis (33 cases), influenza-associated hospitalizations (27), Hepatitis C (15 cases), and gonorrhea (5 cases). Chlamydia, campylobacteriosis, influenza-associated hospitalizations, and Hepatitis C were also in the top five diseases reported during 2015 and 2016.


Table 1. on Page 4 illustrates all of the diseases reported in the community and the number of cases for each of these illnesses. The remainder of this document provides epidemiological data on each of these illnesses as well as brief demographic information on the cases and disease trends over the past five years.

ANNUAL COMMUNICABLE DISEASES

Table 1. Communicable Diseases Reported in Wyandot County, 2017

Class B Reportable Diseases	
Babesiosis	1
Campylobacteriosis	33
Chlamydia infection	57
Cryptosporidiosis	1
<i>E. coli</i>	2
Gonorrhea	5
<i>Haemophilus influenza</i>	1
Hepatitis B	3
Hepatitis C	15
Influenza-associated hospitalization	27
Lyme Disease	3
Pertussis	1
Salmonellosis	3
Shigellosis	1
Streptococcal—Group A—invasive	1
<i>Streptococcus pneumoniae</i>	3
Total	157
Class C Reportable Diseases—Outbreaks	
Influenza	1
Total	1

ANNUAL COMMUNICABLE DISEASES




Notes:

- Case counts include confirmed, probable, and suspect disease classifications
- Sexually transmitted infections include chlamydia and gonorrhea
- Vaccine preventable illnesses include *Haemophilus influenzae*, influenza-associated hospitalizations, *Streptococcus pneumoniae*, and Pertussis
- Bloodborne pathogens includes Hepatitis C
- Enteric illnesses include campylobacteriosis, cryptosporidiosis, *E. coli*, Salmonella, and shigella
- Vectorborne illness include Lyme disease and babesiosis
- Other illness includes streptococcal disease

CHLAMYDIA

DEMOGRAPHICS


Number of Cases:	57
Average Age:	23.7
Median Age:	22
Age Range:	15-55 years
Female:	66.7%
Male:	33.3%
Increase from 2016:	32.6%



EPIDEMIOLOGY

Infectious Agent:	<i>Chlamydia trachomatis</i> bacteria
Case Definition:	Isolation of <i>Chlamydia trachomatis</i> from a clinical specimen
Symptoms:	Men may suffer from painful urination, urinary frequency, and penile discharge; while women may experience vaginal discharge
Source:	Humans
Mode of Transmission:	Sexually transmitted
Incubation Period:	7-21 days
Prevention:	Abstinence, condom use, and identification and treatment of sexual contacts of those proven to be or suspected of being infected with <i>Chlamydia trachomatis</i>


FIVE YEAR TREND



CAMPYLOBACTERIOSIS

DEMOGRAPHICS


Number of Cases:	33
Average Age:	45.6
Median Age:	49
Age Range:	3 months-88 years
Female:	54.5%
Male:	45.5%
Increase from 2016:	83.3%



EPIDEMIOLOGY

Infectious Agent:	<i>Campylobacter</i> organisms, most commonly <i>Campylobacter jejuni</i>
Case Definition:	Isolation or detection of <i>Campylobacter</i> species with or without diarrhea
Symptoms:	Diarrhea (frequently bloody), abdominal cramps, malaise, fever, headache, nausea, and vomiting
Source:	Poultry, cattle, puppies, kittens, swine, sheep rodents, and birds
Mode of Transmission:	Fecal-orally through undercooked meat, contaminated food or raw milk, or direct contact with infected pets, livestock or infants.
Incubation Period:	1-10 days
Prevention:	Cook meats thoroughly, avoid cross-contamination of foods with raw meat juices, avoid unpasteurized milk and untreated water, and washing hands after contact with animals, bowel movements, changing diapers, and before eating or preparing food.


FIVE YEAR TREND



INFLUENZA-ASSOCIATED HOSPITALIZATIONS

DEMOGRAPHICS


Number of Cases:	27
Average Age:	67.9
Median Age:	74
Age Range:	2 months-93 years
Female:	51.9%
Male:	48.1%
Increase from 2016:	440.0%



EPIDEMIOLOGY

Infectious Agent:	Two main types of Influenza virus: Influenza A and Influenza B; both types include different strains that tend to change from year to year
Case Definition:	An illness compatible with influenza virus infection that results in hospitalization
Symptoms:	Fever, body aches, headache, malaise, nonproductive cough, sore throat, and runny nose
Source:	Humans
Mode of Transmission:	Direct person-to-person contact through droplet spread or via articles recently contaminated with nasopharyngeal secretions.
Incubation Period:	1-4 days
Prevention:	The best prevention is annual vaccination; washing hands after sneezing, coughing or using a tissue; cough into sleeve and not into hands


FIVE YEAR TREND



HEPATITIS C

DEMOGRAPHICS


Number of Cases:	15
Average Age:	33.7
Median Age:	29
Age Range:	16-60 years
Female:	40.0%
Male:	60.0%
Decrease from 2016:	21.1%



EPIDEMIOLOGY

Infectious Agent:	Hepatitis C virus
Case Definition:	A positive test for Hepatitis C virus antibodies or detection of the Hepatitis C virus
Symptoms:	Those infected may be asymptomatic; however, some may experience nausea, vomiting, abdominal pain, loss of appetite, dark urine, and/or jaundice
Source:	Human blood
Mode of Transmission:	Injection drug use through the sharing of needles and other drug paraphernalia contaminated with infected blood; non-professional tattooing or in-home tattooing through shared needles or contaminated equipment; sexual transmission inefficiently spreads the virus (rare)
Incubation Period:	2 weeks—6 months
Prevention:	No vaccine is available


FIVE YEAR TREND



GONORRHEA

DEMOGRAPHICS


Number of Cases:	5
Average Age:	32.8
Median Age:	27
Age Range:	19-59 years
Female:	60.0%
Male:	40.0%
Increase from 2016:	66.7%



EPIDEMIOLOGY

Infectious Agent:	<i>Neisseria gonorrhoeae</i> bacteria
Case Definition:	Isolation of <i>Neisseria gonorrhoeae</i> from a clinical specimen
Symptoms:	Men suffer from painful, frequent urination, and penile discharge; women may experience vaginal discharge, painful urination, and vaginal bleeding between menstrual cycles
Source:	Humans
Mode of Transmission:	Sexually transmitted
Incubation Period:	3-8 days
Prevention:	Abstinence, condom use, and identification and treatment of sexual contacts of those proven to be or suspected of being infected with <i>Neisseria gonorrhoeae</i>

FIVE YEAR TREND



TIMELINESS OF REPORTING

Timely reporting of infectious diseases is important in identifying potential outbreaks and in reducing disease burden. Public health relies on health care providers and laboratories for identification and prompt reporting of these infectious diseases. Timeliness requirements for each reportable disease is dependent of the infectious nature and severity of the disease.

Reporting lag is defined as the difference between the date the case was reported to the local health department and the date of diagnosis. For Class A diseases, median and mean lag time values should be less than 1 since these illnesses are required to be reported to the health department immediately, and for Class B and C diseases, mean and median lag time values should be less than 2 since these illness should be reported to the health department by the end of the next business day.

Table 2 illustrates the lag time for select reportable diseases reported in Wyandot County during 2017.

Table 2. Reporting Lag Time for Selected Reportable Diseases in Wyandot County, 2017

Reportable Disease	Reporting Requirement	Cases (N)	Median (Days)	Mean (Days)
Campylobacteriosis	By end of next business day	33	1.5	2.3
Cryptosporidiosis	By end of next business day	1	6	6
E. coli O157:H7	By end of next business day	2	5	5
Giardia	By end of next business day	0	N/A	N/A
Influenza-Associated Hospitalization	By end of next business day	27	1	2.2
Legionnaires' Disease	By end of next business day	0	N/A	N/A
Pertussis	By end of next business day	1	0	0
Salmonella	By end of next business day	3	2	4.7
Shigella	By end of next business day	1	1	1

Note: Reporting lag time is the difference between the date the case was reported to the local health department and the case's date of diagnosis

Date of diagnosis defaulted to lab specimen collection date or illness onset date if blank

Wyandot County



Public Health

Prevent. Promote. Protect.

127-A S. Sandusky Street
Upper Sandusky, Ohio 43351
(419) 294-3852
<http://www.wyandotohealth.com>

CONTACT

MARY SALIMBENE MERRIMAN, MPH

EPIDEMIOLOGIST

Tel: 937-645-2062

Fax 937-642-1568

mary.merriman@uchd.net



ZACH COLLES, MPH

EPIDEMIOLOGIST

Tel: 937-645-2056

Fax 937-642-1568

zach.colles@uchd.net

Compiled and prepared by:

Mary E. Salimbene Merriman, MPH-Epidemiologist



Division of Health Promotion and Planning

940 London Avenue

Suite 1100

Marysville, Ohio 43040

(937) 642-2053